



The University of New Mexico

Geometry

Ed Angel

Professor of Computer Science,
Electrical and Computer
Engineering, and Media Arts
University of New Mexico



The University of New Mexico

Objectives

- Introduce the elements of geometry
 - Scalars
 - Vectors
 - Points
- Develop mathematical operations among them in a coordinate-free manner
- Define basic primitives
 - Line segments
 - Polygons



The University of New Mexico

Basic Elements

- Geometry is the study of the relationships among objects in an n-dimensional space
 - In computer graphics, we are interested in objects that exist in three dimensions
- Want a minimum set of primitives from which we can build more sophisticated objects
- We will need three basic elements
 - Scalars
 - Vectors
 - Points



Coordinate-Free Geometry

- When we learned simple geometry, most of us started with a Cartesian approach
 - Points were at locations in space $\mathbf{p}=(x,y,z)$
 - We derived results by algebraic manipulations involving these coordinates
- This approach was nonphysical
 - Physically, points exist regardless of the location of an arbitrary coordinate system
 - Most geometric results are independent of the coordinate system
 - Example Euclidean geometry: two triangles are identical if two corresponding sides and the angle between them are identical



The University of New Mexico

Scalars


-
- Need three basic elements in geometry
 - Scalars, Vectors, Points
 - Scalars can be defined as members of sets which can be combined by two operations (addition and multiplication) obeying some fundamental axioms (associativity, commutivity, inverses)
 - Examples include the real and complex number systems under the ordinary rules with which we are familiar
 - Scalars alone have no geometric properties



The University of New Mexico

Vectors

- Physical definition: a vector is a quantity with two attributes
 - Direction
 - Magnitude
- Examples include
 - Force
 - Velocity
 - Directed line segments
 - Most important example for graphics
 - Can map to other types









The University of New Mexico

Vector Operations

- Every vector has an inverse
 - Same magnitude but points in opposite direction
- Every vector can be multiplied by a scalar
- There is a zero vector
 - Zero magnitude, undefined orientation
- The sum of any two vectors is a vector
 - Use head-to-tail axiom





The University of New Mexico

Linear Vector Spaces

- Mathematical system for manipulating vectors
- Operations
 - Scalar-vector multiplication $u = \alpha v$
 - Vector-vector addition: $w = u + v$
- Expressions such as
$$v = u + 2w - 3r$$


Make sense in a vector space



The University of New Mexico

Vectors Lack Position

- These vectors are identical
 - Same length and magnitude




- Vectors spaces insufficient for geometry
 - Need points



The University of New Mexico

Points

- Location in space
- Operations allowed between points and vectors
 - Point-point subtraction yields a vector
 - Equivalent to point-vector addition





The University of New Mexico


Affine Spaces

-
- Point + a vector space
 - Operations
 - Vector-vector addition
 - Scalar-vector multiplication
 - Point-vector addition
 - Scalar-scalar operations
 - For any point define
 - $1 \cdot P = P$
 - $0 \cdot P = \mathbf{0}$ (zero vector)



Lines

- Consider all points of the form
 - $P(\alpha) = P_0 + \alpha \mathbf{d}$
 - Set of all points that pass through P_0 in the direction of the vector \mathbf{d}





Parametric Form

- This form is known as the parametric form of the line
 - More robust and general than other forms
 - Extends to curves and surfaces
- Two-dimensional forms
 - Explicit: $y = mx + h$
 - Implicit: $ax + by + c = 0$
 - Parametric:
$$x(\alpha) = \alpha x_0 + (1-\alpha)x_1$$
$$y(\alpha) = \alpha y_0 + (1-\alpha)y_1$$



The University of New Mexico

Rays and Line Segments


- If $\alpha \geq 0$, then $P(\alpha)$ is the *ray* leaving P_0 in the direction \mathbf{d}

If we use two points to define \mathbf{v} , then

$$P(\alpha) = Q + \alpha (R - Q) = Q + \alpha \mathbf{v}$$

$$= \alpha R + (1 - \alpha)Q$$

For $0 \leq \alpha \leq 1$ we get all the points on the *line segment* joining R and Q







The University of New Mexico

Convexity

- An object is *convex* iff for any two points in the object all points on the line segment between these points are also in the object



convex



not convex



Affine Sums

- Consider the “sum”

$$P = \alpha_1 P_1 + \alpha_2 P_2 + \dots + \alpha_n P_n$$

Can show by induction that this sum makes sense iff

$$\alpha_1 + \alpha_2 + \dots + \alpha_n = 1$$

in which case we have the *affine sum* of the points P_1, P_2, \dots, P_n


- If, in addition, $\alpha_i \geq 0$, we have the *convex hull* of P_1, P_2, \dots, P_n



The University of New Mexico

Convex Hull

- Smallest convex object containing P_1, P_2, \dots, P_n
- Formed by “shrink wrapping” points







The University of New Mexico

Curves and Surfaces

- Curves are one parameter entities of the form $P(\alpha)$ where the function is nonlinear
- Surfaces are formed from two-parameter functions $P(\alpha, \beta)$
 - Linear functions give planes and polygons



$P(\alpha)$




$P(\alpha, \beta)$




The University of New Mexico

Planes

- A plane can be defined by a point and two vectors or by three points



$$P(\alpha, \beta) = R + \alpha u + \beta v$$




$$P(\alpha, \beta) = R + \alpha(Q - R) + \beta(P - R)$$



The University of New Mexico

Triangles



for $0 \leq \alpha, \beta \leq 1$, we get all points in triangle



Normals

- Every plane has a vector n normal (perpendicular, orthogonal) to it
- From point-two vector form $P(\alpha, \beta) = R + \alpha u + \beta v$, we know we can use the cross product to find $n = u \times v$ and the equivalent form

$$(P(\alpha) - P) \cdot n = 0$$

