



The University of New Mexico

Transformations

Ed Angel

Professor of Computer Science,
Electrical and Computer
Engineering, and Media Arts
University of New Mexico



The University of New Mexico

Objectives


-
- Introduce standard transformations
 - Rotation
 - Translation
 - Scaling
 - Shear
 - Derive homogeneous coordinate transformation matrices
 - Learn to build arbitrary transformation matrices from simple transformations



The University of New Mexico

General Transformations

A transformation maps points to other points and/or vectors to other vectors






The University of New Mexico

Affine Transformations

- Line preserving
- Characteristic of many physically important transformations
 - Rigid body transformations: rotation, translation
 - Scaling, shear
- Importance in graphics is that we need only transform endpoints of line segments and let implementation draw line segment between the transformed endpoints



Pipeline Implementation





The University of New Mexico

Notation

We will be working with both coordinate-free representations of transformations and representations within a particular frame

P, Q, R : points in an affine space

u, v, w : vectors in an affine space

α, β, γ : scalars

$\mathbf{p}, \mathbf{q}, \mathbf{r}$: representations of points

-array of 4 scalars in homogeneous coordinates

$\mathbf{u}, \mathbf{v}, \mathbf{w}$: representations of points


-array of 4 scalars in homogeneous coordinates



The University of New Mexico

Translation

- Move (translate, displace) a point to a new location




- Displacement determined by a vector d
 - Three degrees of freedom
 - $P' = P + d$




How many ways?

Although we can move a point to a new location in infinite ways, when we move many points there is usually only one way



object



translation: every point displaced by same vector



Translation Using Representations

Using the homogeneous coordinate representation in some frame

$$\mathbf{p} = [x \ y \ z \ 1]^T$$

$$\mathbf{p}' = [x' \ y' \ z' \ 1]^T$$

$$\mathbf{d} = [dx \ dy \ dz \ 0]^T$$

Hence $\mathbf{p}' = \mathbf{p} + \mathbf{d}$ or

$$x' = x + d_x$$

$$y' = y + d_y$$

$$z' = z + d_z$$

note that this expression is in four dimensions and expresses point = vector + point



Translation Matrix

We can also express translation using a 4 x 4 matrix \mathbf{T} in homogeneous coordinates $\mathbf{p}' = \mathbf{T}\mathbf{p}$ where


$$\mathbf{T} = \mathbf{T}(d_x, d_y, d_z) = \begin{bmatrix} 1 & 0 & 0 & d_x \\ 0 & 1 & 0 & d_y \\ 0 & 0 & 1 & d_z \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

This form is better for implementation because all affine transformations can be expressed this way and multiple transformations can be concatenated together



Rotation (2D)

Consider rotation about the origin by θ degrees
- radius stays the same, angle increases by θ





The University of New Mexico

Rotation about the z axis

- Rotation about z axis in three dimensions leaves all points with the same z
 - Equivalent to rotation in two dimensions in planes of constant z

$$x' = x \cos \theta - y \sin \theta$$

$$y' = x \sin \theta + y \cos \theta$$

$$z' = z$$

- or in homogeneous coordinates

$$\mathbf{p}' = \mathbf{R}_z(\theta)\mathbf{p}$$



The University of New Mexico

Rotation Matrix

$$\mathbf{R} = \mathbf{R}_Z(\theta) = \begin{bmatrix} \cos \theta & -\sin \theta & 0 & 0 \\ \sin \theta & \cos \theta & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$



Rotation about x and y axes

- Same argument as for rotation about z axis
 - For rotation about x axis, x is unchanged
 - For rotation about y axis, y is unchanged

$$\mathbf{R} = \mathbf{R}_x(\theta) = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & \cos \theta & -\sin \theta & 0 \\ 0 & \sin \theta & \cos \theta & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

$$\mathbf{R} = \mathbf{R}_y(\theta) = \begin{bmatrix} \cos \theta & 0 & \sin \theta & 0 \\ 0 & 1 & 0 & 0 \\ -\sin \theta & 0 & \cos \theta & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$



The University of New Mexico

Scaling

Expand or contract along each axis (fixed point of origin)


$$x' = s_x x$$

$$y' = s_y y$$

$$z' = s_z z$$

$$\mathbf{p}' = \mathbf{S}\mathbf{p}$$

$$\mathbf{S} = \mathbf{S}(s_x, s_y, s_z) = \begin{bmatrix} s_x & 0 & 0 & 0 \\ 0 & s_y & 0 & 0 \\ 0 & 0 & s_z & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$






The University of New Mexico

Reflection

corresponds to negative scale factors





Inverses

- Although we could compute inverse matrices by general formulas, we can use simple geometric observations
 - Translation: $\mathbf{T}^{-1}(d_x, d_y, d_z) = \mathbf{T}(-d_x, -d_y, -d_z)$
 - Rotation: $\mathbf{R}^{-1}(\theta) = \mathbf{R}(-\theta)$
 - Holds for any rotation matrix
 - Note that since $\cos(-\theta) = \cos(\theta)$ and $\sin(-\theta) = -\sin(\theta)$
 $\mathbf{R}^{-1}(\theta) = \mathbf{R}^T(\theta)$
 - Scaling: $\mathbf{S}^{-1}(s_x, s_y, s_z) = \mathbf{S}(1/s_x, 1/s_y, 1/s_z)$



Concatenation

- We can form arbitrary affine transformation matrices by multiplying together rotation, translation, and scaling matrices
- Because the same transformation is applied to many vertices, the cost of forming a matrix $\mathbf{M}=\mathbf{ABCD}$ is not significant compared to the cost of computing \mathbf{Mp} for many vertices \mathbf{p}
- The difficult part is how to form a desired transformation from the specifications in the application



Order of Transformations

- Note that matrix on the right is the first applied
- Mathematically, the following are equivalent

$$\mathbf{p}' = \mathbf{A}\mathbf{B}\mathbf{C}\mathbf{p} = \mathbf{A}(\mathbf{B}(\mathbf{C}\mathbf{p}))$$

- Note many references use column matrices to represent points. In terms of column matrices

$$\mathbf{p}'^T = \mathbf{p}^T \mathbf{C}^T \mathbf{B}^T \mathbf{A}^T$$



General Rotation About the Origin


A rotation by θ about an arbitrary axis can be decomposed into the concatenation of rotations about the x , y , and z axes

$$\mathbf{R}(\theta) = \mathbf{R}_z(\theta_z) \mathbf{R}_y(\theta_y) \mathbf{R}_x(\theta_x)$$

θ_x θ_y θ_z are called the Euler angles

Note that rotations do not commute

We can use rotations in another order but with different angles






Rotation About a Fixed Point other than the Origin

Move fixed point to origin

Rotate

Move fixed point back

$$\mathbf{M} = \mathbf{T}(p_f) \mathbf{R}(\theta) \mathbf{T}(-p_f)$$






Instancing

- In modeling, we often start with a simple object centered at the origin, oriented with the axis, and at a standard size
- We apply an *instance transformation* to its vertices to

Scale
Orient
Locate






The University of New Mexico

Shear

- Helpful to add one more basic transformation
- Equivalent to pulling faces in opposite directions





Shear Matrix

Consider simple shear along x axis

$$x' = x + y \cot \theta$$

$$y' = y$$

$$z' = z$$

$$\mathbf{H}(\theta) = \begin{bmatrix} 1 & \cot \theta & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

