



The University of New Mexico

---

# Shading II

Ed Angel

Professor of Computer Science,  
Electrical and Computer  
Engineering, and Media Arts  
University of New Mexico



The University of New Mexico

# Objectives

- 
- Continue discussion of shading
  - Introduce modified Phong model
  - Consider computation of required vectors



The University of New Mexico

# Ambient Light

- Ambient light is the result of multiple interactions between (large) light sources and the objects in the environment
- Amount and color depend on both the color of the light(s) and the material properties of the object
- Add  $k_a I_a$  to diffuse and specular terms


reflection coef

intensity of ambient light



# Distance Terms

- The light from a point source that reaches a surface is inversely proportional to the square of the distance between them
- We can add a factor of the form  $1/(ad + bd + cd^2)$  to the diffuse and specular terms
- The constant and linear terms soften the effect of the point source





# Light Sources

---

- In the Phong Model, we add the results from each light source
- Each light source has separate diffuse, specular, and ambient terms to allow for maximum flexibility even though this form does not have a physical justification
- Separate red, green and blue components
- Hence, 9 coefficients for each point source

$$- I_{dr}, I_{dg}, I_{db}, I_{sr}, I_{sg}, I_{sb}, I_{ar}, I_{ag}, I_{ab}$$



The University of New Mexico

# Material Properties

---

- Material properties match light source properties
  - Nine absorption coefficients
    - $k_{dr}$ ,  $k_{dg}$ ,  $k_{db}$ ,  $k_{sr}$ ,  $k_{sg}$ ,  $k_{sb}$ ,  $k_{ar}$ ,  $k_{ag}$ ,  $k_{ab}$
  - Shininess coefficient  $\alpha$




The University of New Mexico

# Adding up the Components

For each light source and each color component, the Phong model can be written (without the distance terms) as

$$I = k_d I_d \mathbf{l} \cdot \mathbf{n} + k_s I_s (\mathbf{v} \cdot \mathbf{r})^\alpha + k_a I_a$$

For each color component we add contributions from all sources





The University of New Mexico

# Modified Phong Model

---

- The specular term in the Phong model is problematic because it requires the calculation of a new reflection vector and view vector for each vertex
- Blinn suggested an approximation using the halfway vector that is more efficient






The University of New Mexico

# The Halfway Vector

- $\mathbf{h}$  is normalized vector halfway between  $\mathbf{l}$  and  $\mathbf{v}$

$$\mathbf{h} = (\mathbf{l} + \mathbf{v}) / |\mathbf{l} + \mathbf{v}|$$





# Using the halfway angle

---


- Replace  $(\mathbf{v} \cdot \mathbf{r})^\alpha$  by  $(\mathbf{n} \cdot \mathbf{h})^\beta$
- $\beta$  is chosen to match shininess
- Note that halfway angle is half of angle between  $\mathbf{r}$  and  $\mathbf{v}$  if vectors are coplanar
- Resulting model is known as the modified Phong or Blinn lighting model
  - Specified in OpenGL standard



The University of New Mexico

# Example

Only differences in these teapots are the parameters in the modified Phong model





The University of New Mexico

# Computation of Vectors

---


- $\mathbf{l}$  and  $\mathbf{v}$  are specified by the application
- Can compute  $\mathbf{r}$  from  $\mathbf{l}$  and  $\mathbf{n}$
- Problem is determining  $\mathbf{n}$
- For simple surfaces  $\mathbf{n}$  can be determined but how we determine  $\mathbf{n}$  differs depending on underlying representation of surface
- OpenGL leaves determination of normal to application
  - Exception for GLU quadrics and Bezier surfaces (Chapter 11)



# Plane Normals

- Equation of plane:  $ax+by+cz+d = 0$
- From Chapter 4 we know that plane is determined by three points  $p_0, p_2, p_3$  or normal  $\mathbf{n}$  and  $p_0$
- Normal can be obtained by

$$\mathbf{n} = (p_2 - p_0) \times (p_1 - p_0)$$






The University of New Mexico

# Normal to Sphere

- Implicit function  $f(x,y,z)=0$
- Normal given by gradient
- Sphere  $f(\mathbf{p})=\mathbf{p}\cdot\mathbf{p}-1$
- $\mathbf{n} = [\partial f/\partial x, \partial f/\partial y, \partial f/\partial z]^T = \mathbf{p}$






# Parametric Form

- For sphere

$$x = x(u, v) = \cos u \sin v$$

$$y = y(u, v) = \cos u \cos v$$

$$z = z(u, v) = \sin u$$



- Tangent plane determined by vectors

$$\frac{\partial \mathbf{p}}{\partial u} = \left[ \frac{\partial x}{\partial u}, \frac{\partial y}{\partial u}, \frac{\partial z}{\partial u} \right]^T$$

$$\frac{\partial \mathbf{p}}{\partial v} = \left[ \frac{\partial x}{\partial v}, \frac{\partial y}{\partial v}, \frac{\partial z}{\partial v} \right]^T$$

- Normal given by cross product

$$\mathbf{n} = \frac{\partial \mathbf{p}}{\partial u} \times \frac{\partial \mathbf{p}}{\partial v}$$



The University of New Mexico

# General Case

---

- We can compute parametric normals for other simple cases
  - Quadrics
  - Parameteric polynomial surfaces
    - Bezier surface patches (Chapter 11)